

Theoretical homework 2

1. Consider a random sample of 25 households. For each household j ($=1, \dots, 25$) you observe their annual expenditures on dining out, Y_j , measured in units of \$1,000, and their annual income, X_j , measured in units of \$10,000. The model that relates Y_j to X_j is

$$\ln(Y_j) = \alpha + \beta \ln(X_j) + U_j, \quad (1)$$

where the errors U_j are normally distributed with zero expectation and variance σ^2 , and are independent of X_j . The OLS estimation results for model (1) are:

$$\begin{array}{rcll} \ln(Y) & = & 0.25 & + & 0.95 \ln(X) & & R^2 = 0.70 \\ & & (0.5) & & (3.8) & & \hat{\sigma}^2 = 0.75 \end{array} \quad (2)$$

where $\hat{\sigma}^2$ is the unbiased estimator of the error variance σ^2 : $E[\hat{\sigma}^2] = \sigma^2$. The numbers between brackets are the t-values.

- (a) If it is known that $E[Y_j|X_j = 10] = 1$ and $E[Y_j|X_j = 5] = 0.5$, what can be said about α and/or β ?
- (b) Test the null hypothesis $\alpha = 0$ against the alternative hypothesis $\alpha \neq 0$ at the 5% significance level.
- (c) If you accept the null hypothesis $\alpha = 0$, would you re-estimate the model without an intercept?
- (d) Test the null hypothesis $\beta = 1$ against the alternative hypothesis $\beta \neq 1$ at the 5% significance level.
- (e) Forecast $\ln(Y)$ for the case $X = 1$. Given that $X_j \neq 1$ for $j = 1, 2, \dots, 25$, show that the standard error of the forecast error equals 1.
- (f) Compute the 95% confidence interval of $\ln(Y)$ for the case $X = 1$.

Value of the distribution function $F(x)$ of the standard normal distribution for x the sum of a value in the first column and the first row.

x	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986

Critical values of the two-sided t test

with m degrees of freedom

% significance levels

m	5%	10%
1	12.704	6.313
2	4.303	2.920
3	3.183	2.353
4	2.776	2.132
5	2.571	2.015
6	2.447	1.943
7	2.365	1.895
8	2.306	1.859
9	2.262	1.833
10	2.228	1.813
11	2.201	1.796
12	2.179	1.782
13	2.160	1.771
14	2.145	1.761
15	2.131	1.753
16	2.120	1.746
17	2.110	1.740
18	2.101	1.734
19	2.093	1.729
20	2.086	1.725
21	2.080	1.721
22	2.074	1.717
23	2.069	1.714
24	2.064	1.711
25	2.059	1.708
26	2.056	1.706
27	2.052	1.703
28	2.048	1.701
29	2.045	1.699
30	2.042	1.697

m = degrees of freedom